

## 5x Hot Start OmniTaq 2 PCR Kit Cat #: HS322



**Amount:** 125  $\mu$ l Enzyme  
2 x 1.25 ml tubes of 5x TM-PCR-Mix  
(sufficient for 500 x 25  $\mu$ l reactions)

**Shipping conditions:** Ice Pack

**Storage conditions:** For best performance, store at -20°C

**Thermostability:** Retains at least 85% activity after 1 hour at 95°C

**Shelf life:** At least 1 year if stored at -20°C and 10 freeze/thaws or at least 1 month if stored at 4°C.

### PRODUCT DESCRIPTION:

Our PCR kit contains 5X concentrated master mix lacking only the Hot Start OmniTaq 2 enzyme (5x TM-PCR-Mix). The enzyme is provided in a separate vial, which allows an adjustment of its final concentration in PCR.

OmniTaq 2 DNA polymerase is a mutant of Taq polymerase that provides 2-3x faster PCR and some inhibition-resistance. It is made with aptamer-based technology, enabling room temperature reaction set-up. *Please note that Hot Start OmniTaq 2 is not yet optimized for use in RT-PCR or RT-LAMP.*

This kit can be used for conventional as well as real-time PCR. For real-time reactions you may need to add a fluorescent dye as an alternative to probes. 5x TM-PCR-Mix composition is: 250 mM Tris-Cl, 80 mM ammonium sulfate, 0.13% Brij 58, 1 mM each dNTP, and 12.5 mM magnesium chloride. Final pH is 9.1

### TYPICAL PCR PROTOCOL for a 25 $\mu$ l reaction:

Reagent	Volume	Final Concentration
5x TM-PCR-Mix	5 $\mu$ l	1x
Left Primer	variable	200 nM
Right Primer	variable	200 nM
DNA template†	variable	0.5-100 ng
PCR Enhancer Cocktail (optional)*	12.5 $\mu$ l	1x
Hot Start OmniTaq 2 enzyme**	0.05 – 0.25 $\mu$ l **	
De-ionized distilled H2O	Adjust final volume to 25 $\mu$ l	

† DNA amount depends mostly on genome size and target gene copy number.

\* If inhibition-resistance is needed, we recommend using one of our PCR Enhancer Cocktails (PEC-1, PEC-1GC, PEC-2, or PEC-2-GC) which are specially formulated for use with whole blood, serum or plasma.

\*\* To determine specific optimal enzyme concentration, we strongly recommend an enzyme titration test for each target. Good starting amount of the enzyme per 25  $\mu$ l reaction is 0.05  $\mu$ l for purified DNA templates and 0.25  $\mu$ l for crude samples containing 5-10% whole blood, plasma or serum. Targets larger than 1 kb may require more enzyme or may benefit from the use of an LA (Long Accurate) version of the polymerase.

### CYCLING CONDITIONS:

1. Initial Denaturing: 94° for 2-8 minutes recommended for crude samples containing 5-10% whole blood, plasma or serum.
2. Denaturing: 94° for 40-60 seconds
3. Annealing: 50°-68° depending on the specific Tm primers for 40-60 seconds
4. Extension: 68° for 1 min/kb target
5. Repeat steps 2-4 for 25-40 cycles

Please visit us on the web at [www.klentaq.com](http://www.klentaq.com) for troubleshooting and detailed protocols.

## REFERENCE:

Barnes, W. M., et al. (2021) A Single Amino Acid Change to Taq DNA Polymerase Enables Faster PCR, Reverse Transcription and Strand-Displacement. *Frontiers in Bioengineering and Biotechnology*. 8:553474.  
doi: 10.3389/fbioe.2020.553474 <https://doi.org/10.3389/fbioe.2020.553474>