5x Klentaq1 PCR Kit Cat #: 120

Amount: 125 µl enyzme and two 1.25 ml tubes of 5x Mastermix

(sufficient for 500 x 25 µl reactions) **Shipping conditions:** Ambient temperature

Storage conditions: -20°C F

Thermostability: Retains at least 85% activity after 1 hour at 95°C

Shelf life: At least 1 year from date of receipt under proper storage conditions.

PRODUCT DESCRIPTION:

Our 5x ready-to-use PCR kit contains Klentaq1, a 5'-exonuclease deficient Taq polymerase with improved fidelity and thermostability. This kit can be used for conventional as well as real-time PCR. For real-time applications you may need to add a fluorescent dye as an alternative to probes. 5x Mastermix composition is: 250 mM Tris-Cl pH 9.2, 80 mM ammonium sulfate, 0.25% Brij 58, 17.5 mM Magnesium Chloride, and 1 mM each dNTP.

TYPICAL PCR PROTOCOL for a 25 ul reaction:

Reagent	Volume	Final Concentration
5x Mastermix	5 μ1	1x
Left Primer	variable	200 nM
Right Primer	variable	200 nM
DNA template†	variable	0.1-100 ng
PCR Enhancer Cocktail (recommended)*	12.5 μ1	1x
Klentaq1	0.05 – 0.25 μl **	
De-ionized distilled H2O	Adjust final volume to 25 ul	

[†] DNA amount depends mostly on genome size and target gene copy number.

CYCLING CONDITIONS:

1. Denaturing: 94° for 2 minutes for 1 cycle

2. Denaturing: 94° for 40-60 seconds

3. Annealing: 50°-68° depending on the specific primers (5° less than Tm) for 40-60 seconds

4. Extension: 68° for 2 min/kb target

5. Repeat steps 2-4 for 25-40 cycles

Please visit us on the web at www.klentaq.com for troubleshooting and detailed protocols.

REFERENCES:

Barnes, W.M. (1994) PCR amplification of up to 35 kb DNA with high fidelity and high yield from bacteriophage templates, PNAS 91, 2216-2220.

Notice to Purchaser: DNA Polymerase Technology products may not be resold, modified for resale or used to manufacture products without an agreement with DNA Polymerase Technology, Inc. The Omni mutant DNA Polymerases are trademarked and patented (US 7,462,475, and US patent pending). No license for OmniTaq 3, to be used in a Polymerase Chain Reaction, has been purchased by DNA Polymerase Technology, Inc.

^{*} For optimal performance, we recommend using one of our PCR Enhancer Cocktails (PEC-1, PEC-1GC, PEC-2, or PEC-2-GC) or 1.3 M Betaine, a general PCR enhancer.

^{**} To determine specific optimal enzyme concentration, we strongly recommend an enzyme titration test for each target. Good starting amount of the enzyme per 25 ul reaction is 0.05 ul. Targets larger than 1 kb may require more enzyme or may benefit from the LA (Long-Accurate) version of the enzyme