

## 5x Omni Klentaq LA PCR Kit

Cat #: 370



**Amount:** 125 µl Omni Klentaq LA  
2 x 1.25 ml tubes of 5x KM-PCR-Mix  
(sufficient for 500 x 25 µl reactions)

**Shipping conditions:** Ice Pack

**Storage conditions:** -20°C

**Thermostability:** Retains at least 85% activity after 1 hour at 95°C

**Shelf life:** At least 1 year if stored at -20°C and 10 freeze/thaws or at least 1 month if stored at 4°C

**Expiration:** On tube label

### PRODUCT DESCRIPTION:

Our PCR kit contains 5X concentrated master mix (5x KM-PCR Mix) lacking only the Omni Klentaq LA enzyme. The enzyme is provided in a separate vial, which allows an adjustment of its final concentration in PCR.

Omni Klentaq LA is a mutant of Klentaq DNA polymerase known to be resistant to soil, blood, and other PCR inhibitors. The Long-Accurate feature allows for amplification of longer products with higher fidelity and accuracy. This kit can be used for conventional as well as real-time PCR. For real-time applications you may need to add a fluorescent dye as an alternative to probes. LA enzymes are not recommended for use with dUTP. 5x KM-PCR-Mix composition is: 250 mM Tris-Cl pH 9.2, 80 mM ammonium sulfate, 0.13% Brij 58, 17.5 mM Magnesium Chloride, and 1 mM each dNTP.

### TYPICAL PCR PROTOCOL for a 25 µl reaction:

Reagent	Volume	Final Concentration
5x KM-PCR-Mix	5 µl	1x
Left Primer	variable	200 nM
Right Primer	variable	200 nM
DNA template†	variable	0.1-100 ng
PCR Enhancer Cocktail (recommended)*	12.5 µl	1x
Omni Klentaq LA	0.05 – 0.25 µl **	
De-ionized distilled H <sub>2</sub> O	Adjust final volume to 25 µl	

† DNA amount depends mostly on genome size and target gene copy number.

\* For optimal performance, we recommend using one of our PCR Enhancer Cocktails (PEC-1, PEC-1GC, PEC-2, or PEC-2-GC) which are specially formulated for use with whole blood, serum or plasma or 1.3M Betaine, a generic PCR enhancer.

\*\* To determine specific optimal enzyme concentration, we strongly recommend an enzyme titration test for each target. A good starting amount of the enzyme per 25 µl reaction is 0.05 µl for purified DNA templates and 0.25 µl for crude samples containing 5-10% whole blood, plasma or serum. Targets larger than 1 kb may require more enzyme.

### CYCLING CONDITIONS:

1. Denaturing: 94° for 2-8 minutes for 1 cycle \*
2. Denaturing: 94° for 40-60 seconds
3. Annealing: 50°-68° depending on the specific T<sub>m</sub> primers for 40-60 seconds
4. Extension: 68° for 2 min/kb target
5. Repeat steps 2-4 for 25-40 cycles

Initial 2-8 min heating step is recommended for crude samples containing 5-10% whole blood, plasma or serum.

**Please visit us on the web at [www.klentaq.com](http://www.klentaq.com) for troubleshooting and detailed protocols.**

### REFERENCES:

Kermekchiev, M.B., et al. (2003) Cold-sensitive mutants of Taq DNA polymerase provide a hot start for PCR. Nucl Acids Res. 31, 6139-6147.

Kermekchiev, M.B. et al. (2009) Mutants of Taq DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. Nucl. Acids Res., 37 (5):e40 E pub.