

5x Omni Klentaq LA PCR Kit

Cat #: 370



Amount: 125 μ l Omni Klentaq LA
2 x 1.25 ml tubes of 5x KM-PCR-Mix
(sufficient for 500 x 25 μ l reactions)

Shipping conditions: Ambient

Storage conditions: -20°C

Thermostability: Retains at least 85% activity after 1 hour at 95°C

Shelf life: At least 1 year if stored at -20°C and 10 freeze/thaws or at least 1 month if stored at 4°C

Expiration: On tube label

PRODUCT DESCRIPTION:

Our PCR kit contains 5X concentrated master mix (5x KM-PCR Mix) lacking only the Omni Klentaq LA enzyme. The enzyme is provided in a separate vial, which allows an adjustment of its final concentration in PCR.

Omni Klentaq LA is a mutant of Klentaq DNA polymerase known to be resistant to soil, blood, and other PCR inhibitors. The Long-Accurate feature allows for amplification of longer products with higher fidelity and accuracy. This kit can be used for conventional as well as real-time PCR. For real-time applications you may need to add a fluorescent dye as an alternative to probes. LA enzymes are not recommended for use with dUTP. 5x KM-PCR-Mix composition is: 250 mM Tris-Cl pH 9.2, 80 mM ammonium sulfate, 0.13% Brij 58, 17.5 mM Magnesium Chloride, and 1 mM each dNTP.

TYPICAL PCR PROTOCOL for a 25 μ l reaction:

Reagent	Volume	Final Concentration
5x KM-PCR-Mix	5 μ l	1x
Left Primer	variable	200 nM
Right Primer	variable	200 nM
DNA template†	variable	0.1-100 ng
PCR Enhancer Cocktail (recommended)*	12.5 μ l	1x
Omni Klentaq LA	0.05 – 0.25 μ l **	
De-ionized distilled H ₂ O	Adjust final volume to 25 μ l	

† DNA amount depends mostly on genome size and target gene copy number.

* For optimal performance, we recommend using one of our PCR Enhancer Cocktails (PEC-1, PEC-1GC, PEC-2, or PEC-2-GC) which are specially formulated for use with whole blood, serum or plasma or 1.3M Betaine, a generic PCR enhancer.

** To determine specific optimal enzyme concentration, we strongly recommend an enzyme titration test for each target. A good starting amount of the enzyme per 25 μ l reaction is 0.05 μ l for purified DNA templates and 0.25 μ l for crude samples containing 5-10% whole blood, plasma or serum. Targets larger than 1 kb may require more enzyme.

CYCLING CONDITIONS:

1. Denaturing: 94° for 2-8 minutes for 1 cycle *
2. Denaturing: 94° for 40-60 seconds
3. Annealing: 50°-68° depending on the specific T_m primers for 40-60 seconds
4. Extension: 68° for 2 min/kb target
5. Repeat steps 2-4 for 25-40 cycles

Initial 2-8 min heating step is recommended for crude samples containing 5-10% whole blood, plasma or serum.

Please visit us on the web at www.klentaq.com for troubleshooting and detailed protocols.

REFERENCES:

Kermekchiev, M.B., et al. (2003) Cold-sensitive mutants of Taq DNA polymerase provide a hot start for PCR. Nucl Acids Res. 31, 6139-6147.

Kermekchiev, M.B. et al. (2009) Mutants of Taq DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. Nucl. Acids Res., 37 (5):e40 E pub.