

## Cesium Klentaq C LA

**Amount:** 100 µl (2000 x 25 µl reactions)

**Shipping conditions:** Ambient temperature

**Storage conditions:** -20°C F

**Thermostability:** Retains at least 85% activity after 1 hour at 95°C

**Shelf life:** At least 1 year from date of receipt under proper storage conditions.

**Cat #: 290**



### PRODUCT DESCRIPTION:

Cesium Klentaq C LA is a cold-sensitive mutant of Klentaq1 (5'-exonuclease deficient Taq polymerase with improved fidelity and thermostability) with the Long-and-Accurate feature that allows amplification of longer products with higher fidelity and accuracy. Due to its suppressed activity at low temperatures this enzyme is designed for hot start PCR performance. LA enzymes are not recommended for use with dUTP. 10x buffer composition is: 500 mM Tris-Cl pH 9.2, 160 mM ammonium sulfate, 0.25% Brij 58, and 35 mM magnesium chloride.

### TYPICAL PCR PROTOCOL for a 25 ul reaction:

Reagent	Volume	Final Concentration
10x Klentaq mutant reaction buffer	2.5 µl	1x
dNTP mix (10 mM each)	0.5 µl	200 µM each
Left Primer	variable	200 nM
Right Primer	variable	200 nM
DNA template†	variable	0.1-100 ng
PCR Enhancer Cocktail (recommended)*	12.5 µl	1x
Cesium Klentaq C LA	0.05 – 0.25 µl **	
De-ionized distilled H2O	Adjust final volume to 25 ul	

† DNA amount depends mostly on genome size and target gene copy number.

\* For optimal performance, we recommend using one of our PCR Enhancer Cocktails (PEC-1, PEC-1GC, PEC-2, or PEC-2-GC) or 1.3 M Betaine, which is a general PCR enhancer.

\*\* To determine specific optimal enzyme concentration, we strongly recommend an enzyme titration test for each target. Targets larger than 1 kb may require more enzyme.

### CYCLING CONDITIONS:

1. Denaturing: 94° for 2-8 minutes for 1 cycle
2. Denaturing: 94° for 40-60 seconds
3. Annealing: 50°-68° depending on the specific Tm primers for 40-60 seconds
4. Extension: 68° for 2 min/kb target
5. Repeat steps 2-4 for 25-40 cycles

**Please visit us on the web at [www.klentaq.com](http://www.klentaq.com) for troubleshooting and detailed protocols.**

### REFERENCES:

Kermekchiev, M.B., et al. (2003) Cold-sensitive mutants of Taq DNA polymerase provide a hot start for PCR. Nucl Acids Res. 31, 6139-6147.

Kermekchiev, M.B. et al. (2009) Mutants of Taq DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. Nucl. Acids Res., 37 (5):e40 E pub.

**Notice to Purchaser:** DNA Polymerase Technology products may not be resold, modified for resale or used to manufacture products without an agreement with DNA Polymerase Technology, Inc. The Omni mutant DNA Polymerases are trademarked and patented (US 7,462,475, and US patent pending). No license for any enzyme to be used in a Polymerase Chain Reaction, has been purchased by DNA Polymerase Technology, Inc.