

Enzyme Combo

Cat #: 510

Amount:

Klentaq LA: 20 ul (0.05 ul per kb but no more than 0.325 ul/ 25 ul rxn)
OmniTaq LA: 25 ul (0.05- 0.5ul/ 25 ul rxn)
OmniTaq 3 LA: 25 ul (0.05- 0.5ul/ 25 ul rxn)
Omni Klentaq LA: 25 ul (0.05- 0.5ul/ 25 ul rxn)
Omni Klentaq 2 LA: 25 ul (0.05- 0.5ul/ 25 ul rxn)
Cesium Klentaq AC LA: 20 ul (0.05 ul/ 25 ul rxn)
CesiumTaq LA: 20 ul (0.05 ul/ 25 ul rxn)

10x Klentaq Reaction Buffer (1.5 ml) for use with Klentaq LA

10x Klentaq Mutant Reaction Buffer (1.5 ml) for use with Omni Klentaq LA, and Omni Klentaq 2 LA, and Cesium Klentaq AC LA

10x Taq Mutant Reaction Buffer (1.5 ml) for use with OmniTaq LA, OmniTaq 3 LA, and CesiumTaq LA

Shipping conditions: Ambient temperature

Storage conditions: -20°C F

Thermostability: Retains at least 85% activity after 1 hour at 95°C

Shelf life: At least 1 year from date of receipt under proper storage conditions.

PRODUCT DESCRIPTION:

Our Enzyme Combo LA allows you to test 7 of our enzymes to see which one works best for your application. LA enzymes are not recommended for use with dUTP.

10x Klentaq Reaction Buffer is provided for Klentaq LA. Buffer composition is: 500 mM Tris, 160 mM ammonium sulfate, 0.5% Brij 58, and 35 mM magnesium chloride. Final pH is 9.2.

10x Klentaq Mutant Reaction Buffer is provided for Omni Klentaq LA, Omni Klentaq 2 LA, and Cesium Klentaq AC LA. Buffer composition is: 500 mM Tris, 160 mM ammonium sulfate, 0.25% Brij 58, and 35 mM magnesium chloride. Final pH is 9.2.

10x Taq Mutant Reaction Buffer is provided for OmniTaq LA, OmniTaq 3 LA, and CesiumTaq LA. Buffer composition is: 500 mM Tris-Cl, 160 mM ammonium sulfate, 0.25% Brij 58, and 25 mM magnesium chloride. Final pH is 9.1.

We also offer (upon request) 10x reaction buffers at pH 7.9 for better fidelity.

TYPICAL PCR PROTOCOL for a 25 ul reaction:

Reagent	Volume	Final Concentration
10x appropriate buffer (see above)	2.5 µl	1x
dNTP mix (10 mM each)	0.5 µl	200 µM each
Left Primer	variable	200 nM
Right Primer	variable	200 nM
DNA template†	variable	0.1-100 ng
PCR Enhancer Cocktail (recommended)*	12.5 µl	1x
DNA Polymerase	0.05 – 0.25 µl **	
De-ionized distilled H2O	Adjust final volume to 25 ul	

† DNA amount depends mostly on genome size and target gene copy number.

* Our PCR Enhancer Cocktails (sold separately) confer additional inhibition resistance when using whole blood, serum, plasma, soil, and some inhibitory foods.

** To determine specific optimal enzyme concentration, we strongly recommend an enzyme titration test for each target. Optimal concentrations start at 0.05 ul / 25 ul rxn when using purified DNA template. Our Omni enzymes require more enzyme (up to 0.5 µl / 25 ul rxn) for use with crude samples containing 5% or more whole blood, plasma or serum, or crude soil extracts, or food matrices. For all our enzymes, targets larger than 1 kb may require more enzyme.

CYCLING CONDITIONS:

1. Denaturing: 94° for 2-8 minutes for 1 cycle
2. Denaturing: 94° for 40-60 seconds
3. Annealing: 50°-68° depending on the specific T_m primers for 40-60 seconds
4. Extension: 68° for 2 min/kb target
5. Repeat steps 2-4 for 25-40 cycles

Please visit us on the web at www.klentaq.com for troubleshooting and detailed protocols.

REFERENCES:

Kermekchiev, M.B., et al. (2003) Cold-sensitive mutants of Taq DNA polymerase provide a hot start for PCR. Nucl Acids Res. 31, 6139-6147.

Kermekchiev, M.B. et al. (2009) Mutants of Taq DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. Nucl. Acids Res., 37 (5):e40 E pub.

Notice to Purchaser: DNA Polymerase Technology products may not be resold, modified for resale or used to manufacture products without an agreement with DNA Polymerase Technology, Inc. The Omni mutant DNA Polymerases are trademarked and patented (US 7,462,475, and US patent pending). No license for OmniTaq 2, to be used in a Polymerase Chain Reaction, has been purchased by DNA Polymerase Technology, Inc.