
# **Klentaq1** Cat #: 100



**Shipping conditions:** Ambient **Storage conditions:** -20°C

**Thermostability:** Retains at least 85% activity after 1 hour at 95°C

Expiration: On tube label



#### PRODUCT DESCRIPTION:

Klentaq1 is a 5'-exonuclease deficient Taq polymerase (an N-terminal deletion of Taq) with improved fidelity and thermostability. 10x buffer composition is: 500 mM Tris-Cl pH 9.2, 160 mM ammonium sulfate, 0.5% Brij 58, and 35 mM magnesium chloride. We also offer (upon request) 10x buffer at pH 7.9 for better fidelity.

# TYPICAL PCR PROTOCOL for a 25 µl reaction:

Reagent	Volume	Final Concentration
10x Klentaq1 reaction buffer	2.5 μ1	1x
DNTP mix (10 mM)	0.5 ul	200 uM each
Left Primer	variable	200 nM
Right Primer	variable	200 nM
DNA template†	variable	0.1-100 ng
PCR Enhancer Cocktail (recommended)*	12.5 μ1	1x
Klentaq1**	0.05 – 0.25 μ1 **	
De-ionized distilled H <sub>2</sub> O	Adjust final volume to 25 ul	-

<sup>†</sup> DNA amount depends mostly on genome size and target gene copy number.

## **CYCLING CONDITIONS**

1. Denaturing: 94° for 2 minutes for 1 cycle 2. Denaturing: 94° for 20-40 seconds

3. Annealing: 50°-68° depending on the specific primers (5° less than Tm) for 40-60 seconds

4. Extension: 68° for 2 min/kb target

5. Repeat steps 2-4 for 25-40 cycles

Please visit us on the web at www.klentaq.com for troubleshooting and detailed protocols.

### **REFERENCES:**

Barnes, W.M. (1994) PCR amplification of up to 35 kb DNA with high fidelity and high yield from bacteriophage templates, PNAS 91, 2216-2220.

U.S. Patent No. 5,436,149

<sup>\*</sup> For optimal performance, we recommend using one of our PCR Enhancer Cocktails (PEC-1, PEC-1GC, PEC-2, or PEC-2-GC) which are specially formulated for use with whole blood, serum or plasma or 1.3M Betaine, a general PCR enhancer.

<sup>\*\*</sup> To determine specific optimal enzyme concentration, we strongly recommend an enzyme titration test for each target. A good starting amount of the enzyme per 25ul reaction is 0.05ul for purified DNA templates and 0.25ul for crude samples containing 5-10% whole blood, plasma or serum. Targets larger than 1 kb may require more enzyme or may benefit from the LA (Long-Accurate) version of the polymerase.