

Omni Klentaq

Cat #: 340



Amount: 125 µl (500 x 25 µl reactions)

Shipping conditions: Ambient temperature

Storage conditions: -20°C

Thermostability: Retains at least 85% activity after 1 hour at 95°C

Shelf life: At least 1 year from date of receipt under proper storage conditions.

PRODUCT DESCRIPTION:

This multipurpose mutant of Klentaq is able to overcome inhibitors in blood, serum, soil, inhibitory foods, and fluorescent dyes. Omni Klentaq can tolerate 15 - 20% whole blood or serum in the PCR reaction. 10x buffer composition is: 500 mM Tris-Cl pH 9.2, 160 mM ammonium sulfate, 0.25% Brij 58, and 35 mM magnesium chloride.

TYPICAL PCR PROTOCOL for a 25 µl reaction:

Reagent	Volume	Final Concentration
10x Klentaq Mutant Reaction Buffer	2.5 µl	1x
dNTP mix (10 mM each)	0.5 µl	200 µM each
Left Primer	variable	200 nM
Right Primer	variable	200 nM
DNA template†	variable	0.1-100 ng
PCR Enhancer Cocktail (recommended)*	12.5 µl	1x
Omni Klentaq	0.05 – 0.25 µl **	
De-ionized distilled H ₂ O	Adjust final volume to 25 µl	

† DNA amount depends mostly on genome size and target gene copy number.

* For optimal performance, we recommend using one of our PCR Enhancer Cocktails (PEC-1, PEC-1GC, PEC-2, or PEC-2-GC) which are specially formulated for use with whole blood, serum or plasma.

** To determine specific optimal enzyme concentration, we strongly recommend an enzyme titration test for each target. Good starting amount of the enzyme per 25 µl reaction is 0.05 µl for purified DNA templates and 0.25 µl for crude samples containing 5-10% whole blood, plasma or serum. Targets larger than 1 kb may require more enzyme or may benefit from the use of the LA (Long Accurate) version of the polymerase.

CYCLING CONDITIONS:

1. Denaturing: 94° for 2-8 minutes for 1 cycle *
2. Denaturing: 94° for 40-60 seconds
3. Annealing: 50°-68° depending on the specific T_m primers for 40-60 seconds
4. Extension: 68° for 2 min/kb target
5. Repeat steps 2-4 for 25-40 cycles

Initial 2-8 min heating step is recommended for crude samples containing 5-10% whole blood, plasma or serum.

Please visit us on the web at www.klentaq.com for troubleshooting and detailed protocols.

REFERENCES:

Kermekchiev, M.B., et al. (2003) Cold-sensitive mutants of Taq DNA polymerase provide a hot start for PCR. Nucl Acids Res. 31, 6139-6147.

Kermekchiev, M.B. et al. (2009) Mutants of Taq DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. Nucl. Acids Res., 37 (5):e40 E pub.