

PCR Enhancer Cocktail P

Cat #: E650



Amount: 5 x 1.25 ml PEC-P (100 x 25 μ l reactions each)

Shipping conditions: Ambient temperature

Storage conditions: -20° C

Expiration: On tube label. If crystallization occurs, the solutions can be restored by soaking at 50-70°C.

PRODUCT DESCRIPTION:

PCR Enhancer Cocktail P (PEC-P) is particularly recommended for samples containing polyphenolic PCR inhibitors. It has been shown to be effective in overcoming inhibition from plant tissues, bile salts, and fecal samples.

Our family of PEC products are non-betaine based PCR enhancers specifically designed for use with inhibitory templates such as plasma, serum, whole blood, inhibitory food matrices, plant tissue, polyphenolic samples, humic acid, bile, and feces. In many cases amplification may be performed **without DNA purification**. They are highly recommended for use with our inhibition-resistant Taq and Klentaq mutants. When used with a PEC, our enzymes can tolerate at least 25% plasma, serum, or whole blood, and as high as 80% GC content templates. PECs are efficient in conventional or real-time PCR, both in SYBR Green and TaqMan assays. PECs are compatible with most commercially available DNA polymerases, but they are not recommended for use with AmpliTaq Gold.

APPLICATION: Include PEC as one half the volume of your PCR reaction (add 25 μ l to each 50 μ l PCR reaction or 12.5 μ l to each 25 μ l PCR reaction.)

Please visit us on the web at www.klentaq.com for help selecting the appropriate PCR Enhancer Cocktail for your application. We also offer troubleshooting and provide detailed protocols.

REFERENCES:

Zhang, Z., et al. (2010) Direct PCR Amplification of DNA from Crude Samples Using a PCR Enhancer Cocktail and Novel Mutants of Taq. J Mol Diagn, 12 (2): 151-161.